arXiv:0708.0893v1 [math.DG] 7 Aug 2007

UNIFORM SOBOLEV INEQUALITIES FOR MANIFOLDS EVOLVING BY RICCI FLOW

Shu-Yu Hsu

Department of Mathematics
National Chung Cheng University
168 University Road, Min-Hsiung
Chia-Yi 621, Taiwan, R.O.C.
e-mail:syhsu@math.ccu.edu.tw

Aug 7, 2007

Abstract. Let M be a compact n-dimensional manifold, $n\ge 2$, with metric $g(t)$ evolving by the Ricci flow $\partial g_{ij}/\partial t=-2R_{ij}$ in $(0,T)$ for some $T\in\mathbb{R}^+\cup\{\infty\}$ with $g(0)=g_0$. Let $\lambda_0(g_0)$ be the first eigenvalue of the operator $-\Delta_{g_0}+\frac{R(g_0)}{4}$ with respect to g_0. We extend a recent result of R. Ye and prove uniform logarithmic Sobolev inequality and uniform Sobolev inequalities along the Ricci flow for any $n\ge 2$ when either $T<\infty$ or $\lambda_0(g_0)>0$. As a consequence we extend Perelman's local κ-noncollapsing result along the Ricci flow for any $n\ge 2$ in terms of upper bound for the scalar curvature when either $T<\infty$ or $\lambda_0(g_0)>0$.

Recently there is a lot of studies on Ricci flow on manifolds because it is an important tool in understanding the geometry of manifolds [H1–3], [Hs1–3], [KL], [MT], [P1], [P2]. On the other hand given any compact n-dimensional manifold M, $n\ge 2$, with a fixed metric g it is known that Sobolev inequalities hold [He]. More specifically for any $q\in[1,n)$ and p satisfying

$$\frac{1}{p}=\frac{1}{q}-\frac{1}{n} \tag{1}$$

there exists a minimal constant $C_{p,q}(M,g)>0$ such that the following holds

$$\left(\int_M|u|^p dV_g\right)^{\frac{1}{p}}\le C_{p,q}(M,g)\left(\left(\int_M|\nabla u|^q\,dV_g\right)^{\frac{1}{q}}+\frac{1}{\text{vol}_g(M)^{\frac{1}{n}}}\left(\int_M|u|^q dV_g\right)^{\frac{1}{q}}\right) \tag{2}$$

1991 *Mathematics Subject Classification.* Primary 58J35, 53C21 Secondary 46E35.

Key words and phrases. Ricci flow, compact manifold, uniform logarithmic Sobolev inequality, uniform Sobolev inequalities, κ-noncollapsing.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$-TEX

for any $u \in W^{1,q}(M,g)$. Note that since the above inequality is scale invariant with respect to the metric g, $C_{p,q}(M,g) = C_{p,q}(M,\lambda g)$ for any $\lambda > 0$. Let M be a n-dimensional compact manifold, $n \geq 2$, with a metric $g(t)$ evolving by the the Ricci flow

$$\frac{\partial}{\partial t} g_{ij} = -2R_{ij} \quad \text{in } M \times (0,T) \tag{3}$$

for some $T \in \mathbb{R}^+ \cup \{\infty\}$ with initial metric $g(0) = g_0$. Because of the importance of Sobolev inequalities in the analysis on manifolds, it is interesting to know whether there exists a constant $C > 0$ in place of $C_{p,q}(M, g(t))$ such that some variants of (2) hold uniformly along the Ricci flow. In [CL] S.C. Chang and P. Lu proved that the Yamabe constant which corresponds to the case $q = 2$, $p = 2n/(n-2)$, and $n \geq 3$ in (1) has a nonnegative derivative at the initial time of the Ricci flow under a technical assumption.

Recently R. Ye [Ye] by using the monotonicity property of the W-functional of Perelman [P1] proved the uniform logarithmic Sobolev inequality for $n \geq 3$ along the Ricci flow. He also proved that when $T < \infty$, then there exist constants $A > 0$, $B > 0$, such that the uniform Sobolev inequality

$$\left(\int_M |u|^{\frac{2n}{n-2}} dV_{g(t)} \right)^{\frac{n-2}{n}} \leq A \int_M \left(|\nabla u|^2 + \frac{R}{4} u^2 \right) dV_{g(t)} + B \int_M u^2 \, dV_{g(t)}$$

holds along the Ricci flow for any $n \geq 3$. Similar result was also proved by R. Ye [Ye] when $\lambda_0(g_0) > 0$ where $\lambda_0(g_0)$ is the first eigenvalue of the operator $-\Delta_{g_0} + \frac{R(g_0)}{4}$ with respect to g_0. As a consequence R. Ye [Ye] also obtain the κ-noncollapsing result in terms of upper bound on the scalar curvature along the Ricci flow for $n \geq 3$ when either $T < \infty$ or $\lambda_0(g_0) > 0$.

In this paper we will use a modification of the technique of R. Ye [Ye], Qi.S. Zhang [Z] and E.B. Davis [D] to prove the uniform logarithmic Sobolev inequality and uniform Sobolev inequalities (8) along the Ricci flow for any $n \geq 2$, $q \in (1,n)$ and p satisfying (1) when either $T < \infty$ or $\lambda_0(g_0) > 0$. As a consequence we extend Perelman's local κ-noncollapsing result along the Ricci flow in terms of upper bound on the scalar curvature for $n \geq 2$ when either $T < \infty$ or $\lambda_0(g_0) > 0$. Note that Perelman's local κ-noncollapsing result is stated in terms of upper bound for $|\text{Rm}|$ [P1] while the κ-noncollapsing result in [Ye] and in this paper is in terms of upper bound on the scalar curvature.

We will assume that $(M, g(t))$, $0 \leq t < T$, is a n-dimensional compact manifold, $n \geq 2$, with metric $g(t)$ evolving by the Ricci flow (3) for the rest of the paper. For any metric g on M, let $R(g)$ be the scalar curvature of (M,g), $R_- = -\min(0,R)$, and $R(x,t) = R(g(t))(x)$. For any $0 \leq t < T$, $r > 0$, $x \in M$, we let $B_t(x,r)$ be the geodesic ball at x with radius r with respect to the metric $g(t)$ and we let $vol_t(A)$ be the volume of any measurable set $A \subset M$ with respect to the metric $g(t)$.

Note that since

$$\frac{\partial}{\partial t} \sqrt{\det(g_{ij}(t))} = -R,$$

$L^p(M, g(t)) = L^p(M, g_0)$ and $W^{1,p}(M, g(t)) = W^{1,p}(M, g_0)$for all $1 \leq p \leq \infty$. Hence we can write $L^p(M)$ and $W^{1,p}(M)$ for $L^p(M, g(t))$ and $W^{1,p}(M, g(t))$ for any $t \in [0,T)$ and

$1 \le p \le \infty$. For any metric g on M, $1 \le p < \infty$, $f \in L^p(M,g)$, let

$$\|f\|_{p,g} = \left(\int_M |f|^p \, dV_g\right)^{\frac{1}{p}}.$$

When there is no ambiguity we will write $\|f\|_p$ for $\|f\|_{p,g}$.

Theorem 1. *Let $n \ge 2$, $q \in [1,n)$ and p satisfy (1). For any $v \in W^{1,q}(M,g)$, $\|v\|_q = 1$, we have*

$$\int_M |v|^q \log v^2 \, dV_g \le \frac{2n}{q} \log(C_{p,q}(M,g)(\|\nabla v\|_q + vol_g(M)^{-\frac{1}{n}})). \tag{4}$$

Moreover for any $u \in W^{1,2}(M,g)$, $\|u\|_2 = 1$,

$$\int_M u^2 \log u^2 \, dV_g \le n \log((C_{\frac{n\mu}{n-\mu},\mu}(M,g)((2/\mu)\|\nabla u\|_2 + vol_g(M)^{-\frac{1}{n}})) \tag{5}$$

for any $1 \le \mu < 2$.

Proof. We will use a modification of the proof of Theorem 3.1 of [Ye] to prove the lemma. Let $u \in W^{1,q}(M,g)$ satisfy $\|u\|_q = 1$. Since $\log x$ is concave function, by Jensen's inequality and (2),

$$\begin{aligned}
&\int_M |u|^q \log |u|^{p-q} \, dV_g \le \log\left(\int_M |u|^{p-q} \cdot |u|^q \, dV_g\right) = \log \|u\|_p^p \\
\Rightarrow \quad &\int_M |u|^q \log u^2 \, dV_g \le \frac{2p}{p-q} \log(C_{p,q}(M,g)(\|\nabla u\|_q + \mathrm{vol}_g(M)^{-\frac{1}{n}} \|u\|_q)) \\
&\qquad = \frac{2n}{q} \log(C_{p,q}(M,g)(\|\nabla u\|_q + \mathrm{vol}_g(M)^{-\frac{1}{n}})).
\end{aligned}$$

We now let $\mu \in [1,2)$. For any $u \in W^{1,2}(M,g)$ satisfying $\|u\|_2 = 1$. Let $v = |u|^{\frac{2}{\mu}}$. Then $\|v\|_\mu = 1$. By the Holder inequality,

$$\begin{aligned}
\int_M |\nabla v|^\mu \, dV_g &= \left(\frac{2}{\mu}\right)^\mu \int_M |u|^{2-\mu} |\nabla u|^\mu \, dV_g \\
&\le \left(\frac{2}{\mu}\right)^\mu \left(\int_M |\nabla u|^2 \, dV_g\right)^{\frac{\mu}{2}} \left(\int_M u^2 \, dV_g\right)^{1-\frac{\mu}{2}} \\
&= \left(\frac{2}{\mu}\right)^\mu \left(\int_M |\nabla u|^2 \, dV_g\right)^{\frac{\mu}{2}} \\
\Rightarrow \quad \left(\int_M |\nabla v|^\mu \, dV_g\right)^{\frac{1}{\mu}} &\le \frac{2}{\mu} \left(\int_M |\nabla u|^2 \, dV_g\right)^{\frac{1}{2}}.
\end{aligned} \tag{6}$$

Hence $v \in W^{1,\mu}(M,g)$. By putting $q = \mu$, $p = n\mu/(n-\mu)$ and $v = |u|^{\frac{2}{\mu}}$ in (4) and using (6) we get (5) and the lemma follows.

By an argument similar to that of Section 3 of [Ye] but with Theorem 1 replacing Theorem 3.1 of [Ye] in the proof there we get that Theorem 3.3 and Theorem 3.5 of [Ye] remain valid for any $n \geq 2$ with the constants $C_S(M, g)$ and $vol_g(M)^{-1/n}$ there being replaced by $2C_{\frac{n\mu}{n-\mu},\mu}(M, g)/\mu$ and $C_{\frac{n\mu}{n-\mu},\mu}(M, g)\mathrm{vol}_g(M)^{-\frac{1}{n}}$ respectively for any $1 \leq \mu < 2$. Hence Theorem A, Theorem B, Theorem C, and the Corollary on P.1–3 of [Ye] remain valid for any $n \geq 2$ with the constants $C_S(M, g_0)$ and $vol_{g_0}(M)^{-1/n}$ there being replaced by $2C_{\frac{n\mu}{n-\mu},\mu}(M, g_0)/\mu$ and $C_{\frac{n\mu}{n-\mu},\mu}(M, g_0)\mathrm{vol}_{g_0}(M)^{-\frac{1}{n}}$ respectively for any $1 \leq \mu < 2$. In particular we have

Theorem 2. *(cf. Theorem A and Theorem C of* [Ye]*) Let M be a compact n-dimensional manifold, $n \geq 2$, with metric $g(t)$ evolving by the Ricci flow (2) in $(0, T)$ for some $T \in \mathbb{R}^+ \cup \{\infty\}$ and initial metric $g(0) = g_0$. Then there exist constants $C_1 > 0$, $C_2 > 0$, depending only on n, $C_{\frac{n}{n-1},1}(M, g_0)$, the upper bound of $vol_{g_0}(M)^{-1}$ and $\max_M R(g_0)_-$ such that*

$$\int_M u^2 \log u^2 \, dV_{g(t)} \leq \sigma \int_M \left(|\nabla u| + \frac{R}{4} u^2 \right) dV_{g(t)} - \frac{n}{2} \log \sigma + C_1 (t + \frac{\sigma}{4}) + C_2$$

holds for any $\sigma > 0$, $0 \leq t < T$, and $u \in W^{1,2}(M)$ with $\|u\|_{2,g(t)} = 1$.

If $\lambda_0 = \lambda_0(g_0) > 0$, then there exists a constant $C_3 > 0$, depending only on n, $C_{\frac{n}{n-1},1}(M, g_0)$, a positive lower bound for $\lambda_0 = \lambda_0(g_0)$, the upper bound of $vol_{g_0}(M)^{-1}$ and $\max_M R(g_0)_-$ such that

$$\int_M u^2 \log u^2 \, dV_{g(t)} \leq \sigma \int_M \left(|\nabla u| + \frac{R}{4} u^2 \right) dV_{g(t)} - \frac{n}{2} \log \sigma + C_3$$

holds for any $\sigma > 0$, $0 \leq t < T$, and $u \in W^{1,2}(M)$ with $\|u\|_{2,g(t)} = 1$.

For any $0 \leq s < T$, let

$$H_s = -\Delta^s + \frac{(R(g(s)) + \max_M R(g(s))_-)}{4}$$

where Δ^s is the Laplacian of M with respect to the metric $g(s)$. Then H_s is a self-adjoint operator on $L^2(M)$. Let

$$Q_s(u) = \int_M \left(|\nabla u|^2_{g(s)} + \frac{(R(g(s)) + \max_M R(g(s))_-)}{4} u^2 \right) dV_{g(s)} \quad \forall u \in W^{1,2}(M)$$

be the associated quadratic form of H_s. By the maximum principle the operator $e^{-H_s t}$ is positivity-preserving for all $t \geq 0$ and is a contraction on $L^\infty(M)$ for all $t \geq 0$. Hence $e^{-H_s t}$ is a symmetric Markov semigroup (P.22 of [D]). By Theorem 2 and the same argument as the proof of Theorem 2.2.7 and Corollary 2.2.8 of [D] we have the following lemma.

Lemma 3. *Suppose $n \geq 2$ and either $T < \infty$ or $\lambda_0(g_0) > 0$. Then there exists a constant $C_4 > 0$ which depends only on T and the constants C_1, C_2, of Theorem 2 if $T < \infty$ and depends only on the constant C_3 of Theorem 2 if $\lambda_0(g_0) > 0$ such that*

$$\|e^{-H_s t} u\|_\infty \leq C_4 t^{-\frac{n}{4}} \|u\|_{2,g(s)} \quad \forall 0 < t \leq 1, 0 \leq s < T, u \in L^2(M).$$

By duality we have

Lemma 4. *Suppose $n \geq 2$ and either $T < \infty$ or $\lambda_0(g_0) > 0$. Let $C_4 > 0$ be as given in Lemma 3. Then*

$$\|e^{-H_s t}u\|_{2,g(s)} \leq C_4 t^{-\frac{n}{4}}\|u\|_{1,g(s)} \quad \forall 0 < t \leq 1, 0 \leq s < T, u \in L^1(M).$$

Corollary 5. *Suppose $n \geq 2$ and either $T < \infty$ or $\lambda_0(g_0) > 0$. Let $C_4 > 0$ be as given in Lemma 3 and let $C_5 = C_4^2$. Then*

$$\|e^{-H_s t}u\|_{\infty} \leq C_5 t^{-\frac{n}{2}}\|u\|_{1,g(s)} \quad \forall 0 < t \leq 1, 0 \leq s < T, u \in L^1(M). \tag{7}$$

Proof. Since $e^{-H_s t}u = e^{-H_s(t/2)}e^{-H_s(t/2)}u$, by Lemma 3 and Lemma 4 for any $0 < t \leq 1$, $0 \leq s < T$, $u \in L^1(M)$,

$$\|e^{-H_s t}u\|_{\infty} \leq C_4 t^{-\frac{n}{4}}\|e^{-H_s(t/2)}u\|_{2,g(s)} \leq C_4^2 t^{-\frac{n}{2}}\|u\|_{1,g(s)}.$$

Theorem 6. *Suppose $n \geq 2$ and either $T < \infty$ or $\lambda_0(g_0) > 0$. Let $1 < q < n$ and p satisfy (1). Then there exists a constant $A_{p,q} = A_{p,q}(M, g_0) > 0$ which depends only on n, T, $C_{\frac{n}{n-1},1}(M, g_0)$, the upper bound of $vol_{g_0}(M)^{-1}$ and* $\max_M R(g_0)_-$ *if $T < \infty$ and depends only on n, $C_{\frac{n}{n-1},1}(M, g_0)$, the lower bound of $\lambda_0(g_0)$, the upper bound of $vol_{g_0}(M)^{-1}$ and* $\max_M R(g_0)_-$ *if $\lambda_0(g_0) > 0$ such that*

$$\left(\int_M |u|^p dV_{g(t)}\right)^{\frac{1}{p}} \leq A_{p,q}\left(\int_M \left(|\nabla u|^2 + \frac{R + 4 + \max_M R(g_0)_-}{4}u^2\right)^{\frac{q}{2}} dV_{g(t)}\right)^{\frac{1}{q}} \tag{8}$$

holds for any $0 \leq t < T$ and $u \in W^{1,q}(M)$.

Proof. We will use a modification of the proof of Theorem 2.4.2 of [D] (cf. [Ye], [Z]) to prove the theorem. Let $\widetilde{H}_s = H_s + 1$. Then $\widetilde{H}_s$ is also a symmetric Markov semigroup. By Corollary 5,

$$\|e^{-\widetilde{H}_s t}u\|_{\infty} \leq C_5 t^{-\frac{n}{2}}\|u\|_{1,g(s)} \quad \forall 0 < t \leq 1, 0 \leq s < T, u \in L^1(M) \tag{9}$$

for some constant $C_5 > 0$ given by (7). Since $e^{-H_s t}$ is a contraction on $L^\infty(M)$ for all $t \geq 0$ and $e^{-\widetilde{H}_s t}u = e^{1-t} \cdot e^{-H_s(t-1)}e^{-\widetilde{H}_s}u$, by (9),

$$\|e^{-\widetilde{H}_s t}u\|_{\infty} \leq e^{1-t}\|e^{-\widetilde{H}_s}u\|_{\infty} \leq C_5 e^{1-t}\|u\|_{1,g(s)} \quad \forall t \geq 1, 0 \leq s < T, u \in L^1(M). \tag{10}$$

By (9) and (10) there exists a constant $C_6 > 0$ such that

$$\|e^{-\widetilde{H}_s t}u\|_{\infty} \leq C_6 t^{-\frac{n}{2}}\|u\|_{1,g(s)} \quad \forall t > 0, 0 \leq s < T, u \in L^1(M). \tag{11}$$

By (11) and the Riesz-Thorin interpolation theorem [D],

$$\|e^{-\widetilde{H}_s t}u\|_{\infty} \leq C_6^{\frac{1}{q}} t^{-\frac{n}{2q}}\|u\|_{q,g(s)} \quad \forall t > 0, 0 \leq s < T, u \in L^q(M). \tag{12}$$

By (12) and an argument similar to the proof of Theorem 2.4.2 of [D] $\widetilde{H}_s^{-\frac{1}{2}}$ is of weak type (p,q) for any $1 < q < n$ and p satisfying (1). We now choose q_1, q_2 such that $1 < q_1 < q < q_2 < n$ and let p_i satisfy (1) with p, q being replaced by p_i and q_i, $i = 1,2$ respectively. Then $p_1 < p < p_2$. By the above argument $\widetilde{H}_s^{-\frac{1}{2}}$ is of weak type (p_1, q_1) and weak type (p_2, q_2). Hence by the Marcinkiewicz interpolation theorem [S] there exists a constant $C > 0$ such that

$$\|\widetilde{H}_s^{-\frac{1}{2}} u\|_{p,g(s)} \le C\|u\|_{q,g(s)} \quad \forall 0 \le s < T. \tag{13}$$

Since R satisfy [H1],

$$R_t = \Delta R + 2|\mathrm{Ric}|^2 \quad \text{ in } M \times (0,T).$$

Hence $\min_M R(x,t) = -\max_M R_-(x,t)$ is a monotone increasing function of $t \in [0,T)$. Then $\max_M R_-(x,t) \le \max_M R(g_0)_-$ for all $0 \le t < T$. Thus by (13) we get (8) and the theorem follows.

Theorem 7. *(κ-noncollapsing result) Suppose $n \ge 2$ and either $T < \infty$ or $\lambda_0(g_0) > 0$. Let $\rho > 0$. Then for any $x_0 \in M$, $0 \le t < T$, $0 < r \le \rho$, satisfying*

$$R(x,t) \le r^{-2} \quad \forall x \in B_t(x_0, r), \tag{14}$$

the following holds

$$vol_t(B_t(x_0,r)) \ge \kappa r^n \tag{15}$$

where

$$\kappa = \min\left(\frac{1}{(2^{(n+3q)/q}A_{p,q})^n}, \frac{1}{(\sqrt{2}A_{p,q}(1 + (4 + \max_M R(g_0)_-)\rho^2)^{1/2})^n}\right) \tag{16}$$

for any $1 < q < n$ and p satisfying (1) where $A_{p,q} = A_{p,q}(M, g_0)$ is given by Theorem 6.

Proof. We will use a modification of the proof of Lemma 2.2 of [He] and Lemma 6.1 of [Ye] to prove the theorem. Suppose there exists $x_0 \in M$, $0 \le t < T$, $0 < r \le \rho$ satisfying (14) such that (15) does not hold. Then

$$\mathrm{vol}_t(B_t(x_0,r)) < \kappa r^n.$$

Let $\overline{g} = r^{-2} g(t)$. Then

$$\mathrm{vol}_{\overline{g}}(B_{\overline{g}}(x_0,1)) < \kappa \tag{17}$$

and

$$\overline{R} \le 1 \quad \text{ in } B_{\overline{g}}(x_0,1) \tag{18}$$

where $\overline{R}$ is the scalar curvature of $\overline{g}$. Let $q \in (1,n)$ and p satisfy (1). By Theorem 6 there exists a constant $A_{p,q} = A_{p,q}(M, g_0) > 0$ such that (8) holds. By (8),

$$\left(\int_M |u|^p dV_{\overline{g}}\right)^{\frac{1}{p}} \le A_{p,q}\left(\int_M \left(|\nabla u|_{\overline{g}}^2 + \frac{\overline{R} + (4 + \max_M R(g_0)_-)\rho^2}{4} u^2\right)^{\frac{q}{2}} dV_{\overline{g}}\right)^{\frac{1}{q}} \tag{19}$$

holds for any $u \in W^{1,q}(M,\overline{g})$. Then for any $u \in W^{1,q}(M,\overline{g})$ with compact support in $B_{\overline{g}}(x_0,1)$ by (16), (17), (18), (19), and the Holder inequality,

$$\begin{aligned}
&\left(\int_{B_{\overline{g}}(x_0,1)} |u|^p dV_{\overline{g}}\right)^{\frac{1}{p}} \\
\le &A_{p,q}\left(\int_{B_{\overline{g}}(x_0,1)} \left(|\nabla u|_{\overline{g}}^2 + \frac{1+(4+\max_M R(g_0)_-)\rho^2}{4}u^2\right)^{\frac{q}{2}} dV_{\overline{g}}\right)^{\frac{1}{q}} \\
\le &\sqrt{2}A_{p,q}\left(\int_{B_{\overline{g}}(x_0,1)} |\nabla u|_{\overline{g}}^q dV_{\overline{g}}\right)^{\frac{1}{q}} \\
&\quad + \sqrt{2}A_{p,q}\frac{(1+(4+\max_M R(g_0)_-)\rho^2)^{\frac{1}{2}}}{2}\left(\int_{B_{\overline{g}}(x_0,1)} |u|^q\, dV_{\overline{g}}\right)^{\frac{1}{q}} \\
\le &\sqrt{2}A_{p,q}\left(\int_{B_{\overline{g}}(x_0,1)} |\nabla u|_{\overline{g}}^q dV_{\overline{g}}\right)^{\frac{1}{q}} \\
&\quad + \sqrt{2}A_{p,q}\frac{(1+(4+\max_M R(g_0)_-)\rho^2)^{\frac{1}{2}}}{2}\kappa^{\frac{1}{n}}\left(\int_{B_{\overline{g}}(x_0,1)} |u|^p\, dV_{\overline{g}}\right)^{\frac{1}{p}} \\
\le &\sqrt{2}A_{p,q}\left(\int_{B_{\overline{g}}(x_0,1)} |\nabla u|_{\overline{g}}^q dV_{\overline{g}}\right)^{\frac{1}{q}} + \frac{1}{2}\left(\int_{B_{\overline{g}}(x_0,1)} |u|^p\, dV_{\overline{g}}\right)^{\frac{1}{p}}.
\end{aligned}$$

Hence

$$\left(\int_{B_{\overline{g}}(x_0,1)} |u|^p\, dV_{\overline{g}}\right)^{\frac{1}{p}} \le 2\sqrt{2}A_{p,q}\left(\int_{B_{\overline{g}}(x_0,1)} |\nabla u|_{\overline{g}}^q dV_{\overline{g}}\right)^{\frac{1}{q}}. \tag{20}$$

For any $0 < r_1 \le 1$, let $u(x) = r_1 - \text{dist}_{\overline{g}}(x,x_0)$. By the Holder inequality,

$$\frac{r_1}{2}\text{vol}_{\overline{g}}(B_{\overline{g}}(x_0,r_1/2))^{\frac{1}{q}} \le \left(\int_{B_{\overline{g}}(x_0,r_1)} |u|^q\, dV_{\overline{g}}\right)^{\frac{1}{q}} \le \text{vol}_{\overline{g}}(B_{\overline{g}}(x_0,r_1))^{\frac{1}{n}}\left(\int_{B_{\overline{g}}(x_0,1)} |u|^p\, dV_{\overline{g}}\right)^{\frac{1}{p}}. \tag{21}$$

Hence by (20) and (21),

$$\begin{aligned}
&\frac{r_1}{2}\text{vol}_{\overline{g}}(B_{\overline{g}}(x_0,r_1/2))^{\frac{1}{q}} \le 2\sqrt{2}A_{p,q}\text{vol}_{\overline{g}}(B_{\overline{g}}(x_0,r_1))^{\frac{n+q}{nq}} \\
\Rightarrow\quad &\text{vol}_{\overline{g}}(B_{\overline{g}}(x_0,r_1)) \ge \left(\frac{r_1}{4\sqrt{2}A_{p,q}}\right)^{\frac{nq}{n+q}}\text{vol}_{\overline{g}}(B_{\overline{g}}(x_0,r_1/2))^{\frac{n}{n+q}} \quad \forall 0 < r_1 \le 1.
\end{aligned} \tag{22}$$

By (17), (22), and an argument similar to the proof of Lemma 2.2 of [He] and [C],

$$\begin{aligned}
&\text{vol}_{\overline{g}}(B_{\overline{g}}(x_0,r_1)) \ge \left(\frac{1}{2^{(n+(5q/2))/q}A_{p,q}}\right)^n r_1^n \quad \forall 0 < r_1 \le 1 \\
\Rightarrow\quad &\kappa > \text{vol}_{\overline{g}}(B_{\overline{g}}(x_0,1)) \ge \left(\frac{1}{2^{(n+(5q/2))/q}A_{p,q}}\right)^n.
\end{aligned}$$

This contradicts (16). Hence (15) holds and the theorem follows.

References

[C] G. Carron, *Inégalités isopérimétriques de Faber-Krahn et conséquences,Actes de la Tables Ronde de Gómétrie Differentielle (Luminy, 1992), 205–232, Sémin. Congr., 1*, Soc. Math. France, Paris, 1966.

[CL] S.C. Chang and P. Lu, *Evolution of Yamabe constants under Ricci flow*, Annals of Global Analysis and Geometry **31** (2007), 147–153.

[D] E.B. Davis, *Heat kernels and spectral theory*, Cambridge University Press, Cambridge, Britain, 1989.

[H1] R.S. Hamilton, *Three-manifolds with positive Ricci curvature*, J. Differential Geom. **17(2)** (1982), 255–306.

[H2] R.S. Hamilton, *Four-manifolds with positive curvature operator*, J. Differential Geom. **24(2)** (1986), 153–179.

[H3] R.S. Hamilton, *The formation of singularities in the Ricci flow*, Surveys in differential geometry, Vol. II (Cambridge, MA, 1993), 7–136, International Press, Cambridge, MA, 1995.

[He] E. Hebey, *Nonlinear Analysis on Manifolds: Sobolev spaces and inequalities*, Courant lecture notes in mathematics vol. 5, Courant Institute of Mathematical Sciences, New York University, New York, 1999.

[Hs1] S.Y. Hsu, *Global existence and uniqueness of solutions of the Ricci flow equation*, Differential and Integral Equations **14(3)** (2001), 305–320.

[Hs2] S.Y. Hsu, *A simple proof on the non-existence of shrinking breathers for the Ricci flow*, Calculus of Variations and P.D.E. **27(1)** (2006), 59–73.

[Hs3] S.Y. Hsu, *Generalized $\mathcal{L}$-geodesic and monotonicity of the generalized reduced volume in the Ricci flow, http://arXiv.org/abs/math.DG/0608197.*

[KL] B. Kleiner and J. Lott, *Notes on Perelman's papers, http://arXiv.org/abs/math.DG/0605667.*

[MT] J.W. Morgan and G. Tian, *Ricci flow and the Poincaré Conjecture, http://arXiv.org/abs/math.DG /0607607.*

[P1] G. Perelman, *The entropy formula for the Ricci flow and its geometric applications, http://arXiv.org /abs/math.DG/0211159.*

[P2] G. Perelman, *Ricci flow with surgery on three-manifolds, http://arXiv.org/abs/math.DG/0303109.*

[S] E.M. Stein, *Singular integral and differentiability properties of functions*, Princeton University Press, Princeton, N.J., U.S.A., 1970.

[Ye] R. Ye, *The logarithmic Sobolev inequality along the Ricci flow, http://arXiv.org/abs/0707.2424v2.*

[Z] Qi S. Zhang, *An uniform Sobolev inequality under Ricci flow http://arXiv.org/abs/0706.1594.*

E-MAIL ADDRESS:SYHSU@MATH.CCU.EDU.TW